

## **Suicide & Depression Screening Risk Guide - Frequency**

|  | No Risk   | Low Risk   | Moderate Risk  | High Risk  |
|--|---|--|--|--|
| Adult Crisis   | Repeat C-SSRS:  | Repeat C-SSRS:   | Repeat C-SSRS:   | Repeat C-SSRS:   |
| Residential<br>(Initial Contact)                             | 1. Decompensation in symptoms of behavior 2. At Discharge (from pathway and/or program)  Repeat PHQ-9: 1. Discharge (from pathway and/or program) | 1. Once Daily 2. If Decompensation in Symptoms 3. At Discharge (from pathway and/or program)  Repeat PHQ-9: 1. Discharge (from pathway and/or program) | 1. Once Daily 2. If Decompensation in Symptoms 3. At Discharge (from pathway and/or program)  Repeat PHQ-9: 1. Discharge (from pathway and/or program) | 1. Once Daily 2. If Decompensation in Symptoms 3. At Discharge (from pathway and/or program)  Repeat PHQ-9: 1. Discharge (from pathway and/or program) |
| Adult Residential (Initial Contact or next Quarterly Review) | Repeat PHQ-9 & C-SSRS:  1. Decompensation in symptoms of behavior  2. Every Quarterly Review  3. At Discharge (from program)                      | Repeat PHQ-9 & C-SSRS:  1. Every 30 Days 2. Decompensation in symptoms or behavior 3. At Discharge (from pathway and/or program)                       | Repeat PHQ-9 & C-SSRS:  1. Every 7 Days 2. Decompensation in symptoms or behavior 3. At Discharge (from pathway and/or program)                        | Repeat PHQ-9 & C-SSRS:  1. Every 3 days 2. Decompensation in symptoms or behavior 3. At Discharge (from pathway and/or program)                        |
| Adult Outpatient (Initial Contact)                           | Repeat PHQ-9 & C-SSRS:  1. Every Appointment  | Repeat PHQ-9 & C-SSRS:  1. Every 7-10 Days 2. Decompensation in symptoms or behavior 3. At Discharge (from pathway and/or program)                     | Repeat PHQ-9 & C-SSRS:  1. Every 5-7 Days 2. Decompensation in symptoms or behavior 3. At Discharge (from pathway and/or program)                      | Repeat PHQ-9 & C-SSRS:  1. Every 1-3 days 2. Decompensation in symptoms or behavior 3. At Discharge (from pathway and/or program)                      |



## **Suicide & Depression Screening Risk Guide - Frequency**

